

Limits on neutrino masses in SO(10) GUT's*

O. Pisanti[†] and L. Rosa[†]

[†] Dipartimento di Scienze Fisiche, Università di Napoli,
Mostra d'Oltremare, Pad.19, I-80125 Napoli, Italy;
INFN, Sezione di Napoli, I-80125 Napoli, Italy.

e-mail: `pisanti@na.infn.it`
e-mail: `rosa@na.infn.it`

Abstract

Studying Renormalization Group Equations for the four typical SO(10) spontaneous symmetry breaking patterns we find the existence of lower limits on neutrino masses.

SO(10) [1] models can be characterized by the intermediate symmetry group (G') appearing when SO(10) breaks down to $SU(3) \otimes SU(2) \otimes U(1) (\equiv G_{SM})$,

$$SO(10) \xrightarrow{M_X} G' \xrightarrow{M_R} G_{SM} \xrightarrow{M_Z} SU(3)_c \otimes U(1)_Q. \quad (1)$$

M_X and M_R are connected to the proton lifetime [2] and Majorana neutrino masses [3] respectively,

$$\tau_{p \rightarrow e^+ \pi^0} = 1.1 \div 1.4 \cdot 10^{32} \left(\frac{M_X}{10^{15} \text{ GeV}} \right)^4 \text{ years}, \quad m_{\nu_i} = \left(\frac{m_\tau}{m_b} \right)^2 \frac{m_{u_i}^2 g_{2R}(M_R)}{M_R y_i(M_R)}, \quad (2)$$

where u_i is the up quark of the i -th generation, g_{2R} is the $SU(2)_R$ coupling and y_i the Yukawa couplings of the i -th fermion generation to the Higgs responsible for the symmetry breaking at M_R .

*Talk given by L. Rosa at the International School on Nuclear Physics; 19th Course: Neutrinos in Astro, Particle and Nuclear Physics, Erice, September 16-24, 1997.

TABLE 1

G'	$M_X/10^{15} \text{ GeV}$	$M_R/10^{11} \text{ GeV}$	G'	$M_X/10^{15} \text{ GeV}$	$M_R/10^{11} \text{ GeV}$
G_{422D}	0.6	420	G_{422}	30	1.1
G_{3221D}	1.9	0.78	G_{3221}	15	0.26

Table 1: Values of M_X and M_R , $\sin^2 \theta_w = 0.2315 \pm 0.0002$; $\alpha_s = 0.118 \pm 0.003$; $\frac{1}{\alpha} = 127.9 \pm 0.09$ [6]

Depending on the representations used to classify the Higgses and on the direction of minimum chosen by the Higgs potential, we obtain the following four patterns of symmetry breaking [4, 5], which represent the four simplest and realistic minimal SO(10) models:

SO(10)	Higgses	G'
(i)	$\xrightarrow{\sigma}$	$G_{422D} \equiv SU(4)_{PS} \otimes SU(2)_L \otimes SU(2)_R \times D$
(ii)	$\xrightarrow{\Phi_1}$	$G_{3221D} \equiv SU(3)_c \otimes SU(2)_L \otimes SU(2)_R \otimes U(1)_{B-L} \times D$
(iii)	$\xrightarrow{\Phi_2}$	$G_{422} \equiv SU(4)_{PS} \otimes SU(2)_L \otimes SU(2)_R$
(iv)	$\xrightarrow{\Phi_3}$	$G_{3221} \equiv SU(3)_c \otimes SU(2)_L \otimes SU(2)_R \otimes U(1)_{B-L},$

(σ is the 54 second rank symmetric representation while Φ is the 210 fourth rank antisymmetric representation of SO(10), $\Phi_1 = \frac{1}{\sqrt{3}}(\Phi_{1234} + \Phi_{3456} + \Phi_{5612})$, $\Phi_2 = \Phi_{78910}$ and $\Phi_3 = \cos\theta\Phi_1 + \sin\theta\Phi_2$).

M_X and M_R are found by solving the following Renormalization Group Equations (RGE) (at first loop)*:

$$\frac{\pi}{11\alpha(M_Z)} \left(\sin^2 \theta_w(M_Z) - \frac{\alpha}{\alpha_S}(M_Z) \right) = \frac{12}{11} \ln \frac{M_R}{M_Z} + C_1 \ln \frac{M_X}{M_R}, \quad (3)$$

$$\frac{\pi}{6\alpha(M_Z)} \left(\frac{3}{2} - 3 \sin^2 \theta_w(M_Z) - \frac{\alpha}{\alpha_S}(M_Z) \right) = \ln \frac{M_R}{M_Z} + C_2 \ln \frac{M_X}{M_R}, \quad (4)$$

where C_1 and C_2 parametrize the dependence of RGE on G' and on the Higgses [4].

A thorough study of these RGE has been carried in [4], making it possible to show that in the four cases an upper limit on M_R exists, which is, trough (2), a lower limit on m_{ν_i} . In table 1 we report the highest values of M_R obtained in the four cases together with the corresponding value of M_X . We see that model (i) is ruled out by the limit on proton decay [6] $M_X \geq 1.6 \cdot 10^{15} \text{ GeV}$. Model (iv) gives

*We use two 10-dimensional representations so to avoid the unwanted relation $m_t = m_b$.

($g_{2R} \sim y_i$) $m_{\nu_\tau} \geq 200 \text{ eV}$, too high with respect to the Cowsik Mc Clelland limit $\sum_i m_{\nu_i} \leq 100 \text{ eV}$ [7]. Model (ii) can eventually be ruled out by the next running of SuperKamiokande [6]. Model (iii) gives $m_{\nu_\tau} \geq 50 \text{ eV}$, the order of magnitude required for the hot solution to the dark matter problem, while $m_{\nu_\mu} \sim 10^{-3} \text{ eV}$ is of the order required by the MSW solution [8] to solar-neutrino problem.

References

- [1] H. Georgi, *Particles and Fields*, ed. C. E. Carlson (1975); H. Fritsch and P. Minkowski, *Ann. of Phys.* **93** (1975) 183.
- [2] F. Buccella, G. Miele, L. Rosa, P. Santorelli, and T. Tuzi, *Phys. Lett.* **B 233** (1989) 178; P. Langaker, *Inner Space and Outer Space*, Eds. E. Kolb et al., Univ. of Chicago Press, 1986.
- [3] M. Gell-Mann, P. Ramond and R. Slansky, *Supergravity* (North-Holland) 1980; T. Yanagida (O.Sawada et al. KEK) 1979.
- [4] F. Acampora, G. Amelino-Camelia, F. Buccella, O. Pisanti, L. Rosa and T. Tuzi, *Nuovo Cimento* **A 108** (1995) 375; F. Buccella and L. Rosa, *Nucl. Phys. B (Proc. Suppl)* **A 28** (1992) 168.
- [5] D.G. Lee, R.N. Mohapatra, M.K. Parida, and M. Rani, *Phys. Rev.* **D 51** (1995) 229.
- [6] Particle Data Book *Phys. Rev.* **D 54** (1996) 48, and Note on Nucleon Decay in Particle Data Book, *Phys. Rev.* **D 50** (1994) 1673; A. Suzuki, invited talk at the International Workshop on Supersymmetry and Unification of Fundamental Interactions, Ecole Polytechnique, Palaiseau, France May, 15-19 1995, TOHOKU-HEP-95-03.
- [7] R. Cowsik and J. Mc Clelland, *Phys. Rev. Lett.* **29** (1972) 669.
- [8] L. Wolfenstein, *Phys.Rev. D* 17 (1978) 2369; S. P. Mikheyev and A. Y. Smirnov, *Sov.J.Nucl.Phys.* 42 (1985) 913.